


QUESTION (2009:1)

(a) Complete the equations below to show how each species will react with water to form an acidic solution.



(b) (i) Determine the $[\text{H}_3\text{O}^+]$, $[\text{OH}^-]$, and pH in each of the following two solutions. $[\text{H}_3\text{O}^+]$ for HCl has been listed.

0.0376 mol L⁻¹ HCl solution

$[\text{H}_3\text{O}^+]/\text{mol L}^{-1}$	$[\text{OH}^-]/\text{mol L}^{-1}$	pH
0.0376		

2.48×10^{-4} mol L⁻¹ NaOH solution

$[\text{H}_3\text{O}^+]/\text{mol L}^{-1}$	$[\text{OH}^-]/\text{mol L}^{-1}$	pH

(ii) Explain why the concentration of the acid, HCl, is equal to the concentration of the hydronium ion, $[\text{H}_3\text{O}^+]$



(iii) The concentration of the hydronium ion, H_3O^+ , in a 0.0376 mol L⁻¹ solution CH_3COOH is less than 0.0376 mol L⁻¹. Explain why the concentration of the hydronium ion is less than 0.0376 mol L⁻¹.

(c) Conductivity of solutions can be described as being high, low, or having no conductivity. Compare and contrast the conductivity of the three solutions shown below.

0.100 mol L⁻¹ HCl, 0.100 mol L⁻¹ CH_3COOH & 0.100 mol L⁻¹ NaOH

QUESTION (2008:3)

Complete the equations below to show how each species will react with water to form an acidic solution.

**QUESTION (2008:4)**

Determine the $[\text{H}_3\text{O}^+]$, $[\text{OH}^-]$ and pH in each of the following solutions.

(a) 0.00112 mol L⁻¹ HCl solution.

(b) 3.68×10^{-2} mol L⁻¹ NaOH solution.

QUESTION (2008:8)

Aqueous solutions of acids HA and HB both have the same concentration of 0.100 mol L⁻¹. The pH of the solution of acid HA is 3.5 and the pH of the solution of acid HB is 1.8.

(a) (i) Identify which one of these acids is stronger and circle your choice. HA HB

(ii) Discuss the reasons for your choice.

You should include relevant equations in your answer, as well as reference to what is meant by the strength of an acid.

- (b) Describe what is observed when the following two tests are carried out on 5 mL samples of the acids HA and HB.
- (i) Identical small pieces of magnesium ribbon are placed in each acid.
- (ii) Sodium hydroxide solution is slowly added to each acid. The volume of sodium hydroxide solution required to completely react with the acid is measured.
- (c) Discuss the observations in (b) (i) and (ii).
Your answer must include reference to:
- similarities and / or differences in the observations of the tests on each acid
 - equations for reactions.

QUESTION (2007:1)

- (a) The bicarbonate ion, HCO_3^- , can both accept and donate hydrogen ions (protons). Complete the equations below.
- $\text{HCO}_3^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ +$ Reaction A
- $\text{HCO}_3^- + \text{H}_2\text{O} \rightleftharpoons \text{OH}^- +$ Reaction B
- (b) When sodium bicarbonate, NaHCO_3 , dissolves in water the solution is basic. Circle Reaction A or Reaction B to show which reaction predominates. Justify your answer.

QUESTION (2007:2)

- (a) Give the pH of 0.125 mol L^{-1} HCl.
- (b) Calculate the hydroxide ion concentration of NaOH solution at pH 10.2.
- (c) Calculate the pH of 0.124 mol L^{-1} NaOH.

QUESTION (2007:7)

Methyl orange can be used as an acid-base indicator. It is pink in solutions with a pH less than 3 and yellow in solutions with a pH greater than 4.

Four beakers are known to contain one each of:

- 0.1 mol L^{-1} HCl
- 0.01 mol L^{-1} HCl
- distilled water
- 0.1 mol L^{-1} NaOH

- (a) Complete the following table.

	pH	Colour of methyl orange
0.1 mol L^{-1} HCl		
0.01 mol L^{-1} HCl		
distilled water		
0.1 mol L^{-1} NaOH		

- (b) Using only the methyl orange indicator, additional water, test tubes and a measuring cylinder, discuss how a student could identify each of the four solutions.

QUESTION (2006:1)

(a) Complete the table below to show the conjugate acid-base pairs.

Conjugate acid	Conjugate base
NH_4^+	
H_2PO_4^-	
	Cl^-
	HSO_4^-

(b) Circle the ion that can act as both an acid and a base. CH_3COO^- HCO_3^-
Justify your choice.

QUESTION (2006:4)

The table below shows the pH of two acids, HA and HB, each with the same concentration. When these acids react with magnesium metal, hydrogen gas (H_2) is produced. Discuss the reactions of both acids, HA and HB, with magnesium metal when the same volume of each acid is used.

Acid	pH
HA	1.00
HB	4.00

In your answer include:

- species in solution
- rate of reaction
- total volume of gas produced.


QUESTION (2006:7)

(a) Complete the table below to show the hydronium ion concentration, hydroxide ion concentration, and pH for the three solutions shown.

$$K_w = 1.00 \times 10^{-14}$$

Solution	$[\text{H}_3\text{O}^+] / \text{mol L}^{-1}$	$[\text{OH}^-] / \text{mol L}^{-1}$	pH
hydrochloric acid (HCl)	0.0720		
sodium hypochlorite (NaOCl)			11.4
hypochlorous acid (HOCl)		2.24×10^{-11}	

(b) Hypochlorous acid is a weak acid. Complete the equation below to show the reaction of hypochlorous acid with water.



(c) A solution of sodium hypochlorite, NaOCl, is basic. Discuss the above statement, including appropriate chemical equation(s) in your answer.

- (a) Complete the following equations to show the reaction of both hydrochloric acid and propanoic acid with water.
- $$\text{HCl} + \text{H}_2\text{O} \rightarrow$$
- $$\text{CH}_3\text{CH}_2\text{COOH} + \text{H}_2\text{O} \rightleftharpoons$$
- (b) Consider the properties described in the table above. Explain the differences in the conductivity and pH of the two acids. In your explanation include reference to the species present in each solution.

QUESTION (2004:6)

- (a) The table below shows four different solutions. Each solution contains both hydronium ions (H_3O^+) and hydroxide ions (OH^-). Complete the table, showing the relative concentrations of these two ions in solution, by placing H_3O^+ ions and OH^- ions in the appropriate box. The first one has been done for you.

Solution	Concentration of H_3O^+ and OH^- in solution	
	Greater than $1 \times 10^{-7} \text{ mol L}^{-1}$	Less than $1 \times 10^{-7} \text{ mol L}^{-1}$
Na_2CO_3 0.100 mol L^{-1}	OH^-	H_3O^+
HNO_3 0.100 mol L^{-1}		
NaOH 0.100 mol L^{-1}		
NH_4Cl 0.100 mol L^{-1}		

- (b) Calculate the pH of a solution with a hydronium ion concentration, $[\text{H}_3\text{O}^+]$, of $0.0350 \text{ mol L}^{-1}$. State your answer to 3 significant figures.
- (c) If a solution of sodium hydrogen carbonate has a pH of 9.20, calculate the concentration of hydroxide ions, $[\text{OH}^-]$, present in the solution. State your answer to 3 significant figures.
- $$K_w = 1.00 \times 10^{-14}$$