


$$\text{pH} = -\log [\text{H}_3\text{O}^+]$$

$$[\text{H}_3\text{O}^+] = 10^{-\text{pH}}$$

$$\text{pOH} = -\log [\text{OH}^-]$$

$$[\text{OH}^-] = 10^{-\text{pOH}}$$

$$K_w = [\text{H}_3\text{O}^+] [\text{OH}^-] = 10^{-14}$$



Enter  $1.35 \times 10^{-4}$  into calculator as

1.35 EXP (-) 4 OR

1.35 x 10 ^ (-) 4

Weak acids

$$K_a = [\text{H}_3\text{O}^+]^2 / [\text{HA}]$$

or

$$[\text{H}_3\text{O}^+] = \sqrt{K_a \times [\text{HA}]}$$

Weak bases

$$K_b = [\text{OH}^-]^2 / [\text{B}]$$

or

$$[\text{OH}^-] = \sqrt{K_b \times [\text{B}]}$$

Level 2 stuff revisited

Equations for calculating pH of weak acids and bases

Weakly acidic salts eg  $\text{NH}_4^+$  which contains  $\text{NH}_4^+$  ion

$$K_a = [\text{H}_3\text{O}^+]^2 / [\text{salt}]$$

or

$$[\text{H}_3\text{O}^+] = \sqrt{K_a \times [\text{salt}]}$$

Weakly basic salts eg  $\text{CH}_3\text{COONa}$  which contains  $\text{CH}_3\text{COO}^-$  ion

$$K_b = [\text{OH}^-]^2 / [\text{salt}]$$

or

$$[\text{OH}^-] = \sqrt{K_b \times [\text{salt}]}$$

$$\text{pH} + \text{pOH} = 14$$

$$\text{p}K_a + \text{p}K_b = \text{p}K_w = 14$$

$$K_a \times K_b = K_w = 10^{-14}$$

$$K_w = [\text{H}_3\text{O}^+] [\text{OH}^-] = 10^{-14}$$

$$\text{p}K_w = \text{pH} + \text{pOH} = 14$$

$$\text{p}K_a = -\log K_a$$

$$\text{p}K_b = -\log K_b$$

$$\text{p}K_w = -\log K_w$$

$$K_a = 10^{-\text{p}K_a}$$

$$K_b = 10^{-\text{p}K_b}$$

$$K_w = 10^{-\text{p}K_w}$$

Equations for calculating pH of salts that are weakly acidic / basic

Converting things when you need something you weren't given

$$\text{pH} = \text{pKa} + \log \frac{[\text{base}]}{[\text{acid}]}$$

Note: when  $[\text{base}] = [\text{acid}]$ ,  
then  $\text{pH} = \text{pKa}$

**Z** Remember this! **e**  
The base is the one **WITHOUT** the  
proton and the acid is the one  
**WITH** the proton!

**poa**

Buffer made from  $\text{CH}_3\text{COOH}$  and  
 $\text{CH}_3\text{COONa}$

$\text{CH}_3\text{COONa}$  dissolves completely in  
water into  $\text{CH}_3\text{COO}^- (\text{aq})$  and  $\text{Na}^+ (\text{aq})$   
 $\text{CH}_3\text{COONa} + \text{aq} \rightarrow \text{CH}_3\text{COO}^- + \text{Na}^+$

$$\text{pH} = \text{pKa} + \log \frac{[\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]}$$

$\text{pKa} (\text{CH}_3\text{COOH})$  will be used in calc.


Alter  $[\text{CH}_3\text{COO}^-] : [\text{CH}_3\text{COOH}]$  ratio  
to adjust the pH

Buffer equation

Acidic buffer example

Buffer made from  $\text{NH}_3$  and  $\text{NH}_4\text{Cl}$





$\text{NH}_4\text{Cl}$  dissolves completely in water  
into  $\text{NH}_4^+ (\text{aq})$  and  $\text{Cl}^- (\text{aq})$




$$\text{pH} = \text{pKa} + \log \frac{[\text{NH}_3]}{[\text{NH}_4^+]}$$

$\text{pKa} (\text{NH}_4^+)$  will be used in calc.

Alter  $[\text{NH}_3] : [\text{NH}_4^+]$  ratio to adjust  
the pH



Basic salt:



(write + aq instead of +  $\text{H}_2\text{O}$  if dissolving)

Basic buffer example

Equations for acids, bases and  
salts in water