


AS 91165

Demonstrate understanding of the properties of selected organic compounds

Collated Flow Chart Type Questions / types of reaction


(2020:1)

- (d) Complete the following reaction scheme by drawing the structural formulae for the organic compounds A, B and C, identifying reagents X and Y, and reaction types 1, 2, and 3.



(2019:2)

- (a) Complete the following reaction scheme for propene, C_3H_6 , by drawing the structural formulae for the organic compounds A to D, naming compound A and identifying Reagents 1 and 2, including any conditions.




- (b) Explain how you identified the major and minor products (C and D) in the reaction of propene with hydrogen bromide solution, $HBr(aq)$.
- (c) Compare and contrast the reaction that forms compound B to the reverse reaction that forms propene, C_3H_6 , from compound B.

(2019:2)

(b) The conversion of bromoethane to chloroethane requires two steps, with alcohol as an intermediate product.

(i) Use this information to complete the reaction scheme below by drawing the structural formulae of each organic molecule and naming the intermediate alcohol and the reagents required.




(ii) Elaborate on the reaction scheme for this conversion. In your answer, you should identify:

- any conditions needed for each step of the conversion
- the type of reaction occurring for each step of the conversion

(2018:2)

(c) Many organic reactions take more than one step in order to convert from one organic molecule to another. Two steps are required to produce pentan-2-amine from an alcohol. Use the information given to analyse the reactions.

(i) Draw the structural formulae of the compounds, and name the reagents involved in the process, in the boxes.




(ii) Elaborate on the reactions in the scheme above. In your answer you should identify:

- any conditions needed for each step of the conversion
- the names of alcohol X and organic molecule Y
- the type of reaction that is occurring for each step of the conversion.

(2017)


- (a) (i) Complete the following reaction scheme by drawing the structural formulae for the organic compounds A, C, and D, and identifying reagents 2 and 3.




- (ii) Identify the types of reactions that occur to produce compounds A, B, C, D, and E.
- (c) Compounds B and E react together.
- (i) Write a balanced equation for the reaction that occurs between compounds B and E.
- (ii) Identify the type of reaction that occurs between compounds B and E. Justify your answer.
- (d) Explain how compound A from the reaction scheme could be directly converted into compound D.

(2016)

- (a) (i) Complete the following chart by drawing the structural formulae for the organic compounds A, B, and C and identifying reagent X




(2015)



- Complete the scheme above by drawing the structural formulae of the organic compounds A to D.
- Circle the functional group of each of the organic compounds A, B, and C that you have drawn.
- Identify reagents X and Y. Reagent X: Reagent Y

(2014)


A reaction scheme is shown below.




- (a) (i) Explain why **Reaction 1** from the reaction scheme is classified as an addition reaction.
 (ii) Explain why compound **A** is the **major** product for **Reaction 1** shown in the reaction scheme.
- (b) (i) Explain why **Reaction 2** from the reaction scheme, is classified as an elimination reaction.
 (ii) **Reaction 4** is also an elimination reaction. Draw the structural formula of the product formed in **Reaction 4**.
- (c) (i) Draw TWO repeating units of the polymer formed in **Reaction 5**.
 (ii) Compare and contrast the polymer formed in **Reaction 5** to the polymer formed in **Reaction 3**. In your answer you should explain why the polymers formed in these two reactions are different.

(2013)

(a) The flow diagram below shows a reaction scheme for the conversion of but-1-ene into but-2-ene.




- (i) Use the reaction scheme above to complete the following table to show:
- the formula of each reagent (A, B and C), including any necessary conditions
 - the type of reaction occurring.
- (ii) For the following reaction:



Circle the words that describe the product formed. **major product** **minor product**
Explain your answer.

(2012)

But-1-ene is used in the reaction sequence shown below.



- (a) (i) Draw two repeating units of the polymer, B, formed in Reaction 1.
- (ii) Give the name or formula of a suitable reagent in Reaction 4; include any specific conditions required.
- (iii) Give the name or formula of a suitable reagent in Reaction 3; include any specific conditions required.
- (b) Can compound A exist as geometric (cis-trans) isomers?
Justify your answer, including reference to the requirements for geometric (cis-trans) isomers.
- (c) (i) Draw the structural formulae of the organic molecules C and D, formed in Reaction 2.
- (ii) Elaborate on the reaction occurring in Reaction 2. In your answer you should include:
- identification of the major and minor products
 - an explanation of why there are two possible products
 - justification of your placement of the different structures in boxes C and D with reference to the reaction sequence.


(2009) From expired standard

For each of the THREE following reactions:


- Write the name or structural formula of the reactant used.
- State the type of reaction occurring. Choose from the list below.

acid-base, addition, elimination, hydrolysis, substitution


Reaction One



Reaction Two



Reaction Three




(2020:1)

- (d) Compound A $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$
 Compound B $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$
 Compound C $\text{CH}_3\text{CH}_2\text{COO}^-$
 Reagent X $\text{KOH}(\text{aq}) / \text{NaOH}(\text{aq})$
 Reagent Y $\text{H}^+ / \text{MnO}_4^-$ or MnO_4^- or $\text{H}^+ / \text{Cr}_2\text{O}_7^{2-}$
 Reaction type 1: addition
 Reaction type 2: substitution
 Reaction type 3: oxidation

(2019:2)

(a)




- A: Polypropene
 R1: conc sulfuric acid
 R2: conc NH_3 (alc)
 B: propan-2-ol OR propan-1-ol
 C: 2-bromopropane
 D: 1-bromopropane

- (b) This is an addition reaction to an asymmetric alkene. When the HBr is added to propene when the double bond breaks, there are two possible products. The H atom is more likely to bond to the carbon with more hydrogens. In propene, the second carbon has one hydrogen and the first carbon has two hydrogens therefore the H from HBr bonds to the first carbon and the Br bonds to the second carbon making 2-bromopropane the major product. 1-bromopropane is the minor product where Br bonds to the first carbon.
- (c) The reaction that forms compound B is an addition reaction where the double bond is broken to add OH and H to saturate the molecule and form an alcohol. The reverse reaction is the removal of the H and OH to form a double bond in an elimination reaction, forming an unsaturated molecule with a double bond. The elimination reaction uses concentrated sulfuric acid to remove the water whereas the addition reaction uses dilute sulfuric acid to add the water. The reactions are opposite in that one breaks the double bond to increase saturation and one forms a double bond to decrease saturation.


(2019:2)

(b) (i)



- (ii) Both reactions are a substitution. In the first step, KOH is in aqueous solution to enable the OH to be substituted for the bromine atom to form ethanol. In the second step, SOCl_2 / PCl_3 / PCl_5 can be used to substitute the OH for a chlorine atom.

(2018)



Reagent 1: SOCl_2 / PCl_3 / PCl_5 / conc HCl or Lucas Reagent

Reagent 2: $\text{NH}_3(\text{alc})$

- (ii) Both reactions are substitution reactions because one atom or group of atoms is substituted by another. In the first step, the OH group on the alcohol, pentan-2-ol is substituted by a Cl atom to make a chloroalkane, 2-chloropentane. The reagent used is SOCl_2 . To convert the chloroalkane to an amine requires conc NH_3 (alc). This causes the Cl to be substituted by an NH_2 to form the amine. (This is so that the OH group in aqueous ammonia does not get substituted onto the chloroalkane.)

(2017)

- (a) (i) Complete the reaction scheme by drawing the structural formulae for the organic compounds A, C, and D, and identifying reagents 2 and 3.
 Compound A. $\text{CH}_3\text{CH}_2\text{OH}$ Compound C. CH_3CH_3 Compound D. $\text{CH}_3\text{CH}_2\text{Cl}$ Reagent 2: $\text{Cr}_2\text{O}_7^{2-} / \text{H}^+$ or $\text{MnO}_4^- / \text{H}^+$ Reagent 3: H_2 / Ni or $\text{H}_2 / \text{Pt} / 150^\circ\text{C}$.

(ii) Identify the types of reactions that occur to produce compounds A, B, C, D, and E.

Reaction A: addition (hydration) Reaction B: oxidation Reaction C: addition (hydrogenation)


Reaction D: substitution (halogenation) Reaction E: substitution

(c) (i) $\text{CH}_3\text{COOH}(\text{aq}) + \text{CH}_3\text{CH}_2\text{NH}_2(\text{aq}) \rightarrow \text{CH}_3\text{COO}^-(\text{aq}) + \text{CH}_3\text{CH}_2\text{NH}_3^+(\text{aq})$ (or amide condensation reaction)

(ii) The reaction between B and E is an acid-base (neutralisation) reaction. Acid-base reactions involve a proton / H^+ transfer. Protons / H^+ , are released from the carboxylic acid functional group, $-\text{COOH}$, resulting in a salt forming containing the $-\text{COO}^-$ group. The proton / H^+ is accepted by the amine functional group, $-\text{NH}_2$, this forms a salt containing the $-\text{NH}_3^+$ group.


(d) PCl_3 / PCl_5 / SOCl_2 can be used to convert compound A, $\text{CH}_3\text{CH}_2\text{OH}$, an alcohol, to the chloroalkane, $\text{CH}_3\text{CH}_2\text{Cl}$. This is a substitution reaction where the $-\text{OH}$ group in compound A is replaced by a Cl atom from PCl_3 / PCl_5 / SOCl_2 .

(2016)



(2015)

(a)




(b) Functional groups circled.

(c) Reagent X is concentrated sulfuric acid, conc H_2SO_4 , or $\text{c.H}_2\text{SO}_4$. Reagent Y is alcoholic potassium hydroxide, $\text{KOH}(\text{alc})$ or alcoholic sodium hydroxide, $\text{NaOH}(\text{alc})$.

(2012)



(a) (i)



(ii) H_2 (/Pt)

(iii) PCl_3 / PCl_5 / SOCl_2

(b) No; for a molecule to exist as geometric isomers, it must contain a double bond, and each carbon (involved in the double bond) must have two different atoms / groups attached to it. Compound A has a double bond, but the atoms attached to one carbon are both the same (two hydrogen atoms) so it does not form a geometric isomer.




(c) (i) C

D


(ii) C is the major product and D is the minor product.

There are 2 possible products because when the double bond is broken, an H (or -OH) will bond to one C (and a -OH group (or H) will bond with the other C). The product will depend on which (C) the H (or the -OH) bond to.



C must be

since product E is



i.e. both functional groups are on the second carbon atom.

If $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ was C then E would be $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$.

(2009)

Reaction One: Reactant A – $\text{H}_2\text{O}/\text{H}^+$ OR $\text{H}_2\text{O}/\text{acid}$ OR concentrated H_2SO_4 then H_2O OR dilute / aq H_2SO_4
Type – addition

Reaction Two: Reactant B – conc / H_2SO_4 Type – elimination OR Al_2O_3 / broken pottery / steel wool

Reaction Three: Reactant C – bromine / Br_2 Type – substitution